## Characteristics of an Ideal Non-invasive Liver Fibrosis Test



|              | Validated Across<br>Multiple Settings | Applicability in Different<br>Patients/Conditions | Economic Health Value<br>(Cost Effectiveness) |                                      | Access               |                         |
|--------------|---------------------------------------|---|---|--------------------------------------|----------------------|-------------------------|
|              |                                       |   | Add Clinical<br>Value                         | Decrease<br>Unnecessary<br>Referrals | Simple<br>Blood Test | Large<br>Installed Base |
| The ELF Test | <b>✓</b>                              | <b>/</b>  | <b>/</b>                                      | <b>/</b>                             | <b>✓</b>             | <b>/</b>                |

- 1. Day J, et al. J Appl Lab Med. 2019 Mar;3:815-26.
- 2. Parkes J, et al. Gut. 2010;59:1245-51.
- 3. Irvine KM, et al. Liver Int. 2016;36:370-7.
- 4. Rosenberg WM, et al. Gastroenterology. 2004;127:1704-13.
- 5. Nobili, et al. Gastroenterology. 2009;136:160-7.
- 6. Srivastava A, et al. J Hepatol. 2019; Apr 6:S0168-8278(19)30227-2. doi:10.1016/j.jhep.2019.03.033.